

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

09/33609, B1
#56



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 38/04, 38/10, 38/16, 39/395, 45/00, C07K 14/46, 14/435, 16/30, C12N 5/20, G01N 33/53, 33/536</p>	<p>A1</p>	<p>(11) International Publication Number: WO 95/20974 (43) International Publication Date: 10 August 1995 (10.08.95)</p>
<p>(21) International Application Number: PCT/US95/00095 (22) International Filing Date: 5 January 1995 (05.01.95) (30) Priority Data: 08/190,411 1 February 1994 (01.02.94) US (71) Applicants: LUDWIG INSTITUTE FOR CANCER RE- SEARCH [US/US]; 1345 Avenue of the Americas, New York, NY 10105 (US). MEMORIAL SLOAN-KETTERING CANCER CENTER [US/US]; 1275 York Avenue, New York, NY 10021 (US). (72) Inventors: CHEN, Yao-Tseng; Cornell Medical Center, 1315 York Avenue, New York, NY 10021 (US). STOCKERT, Elisabeth; 1275 York Avenue, New York, NY 10021 (US). CHEN, Yachi; 1275 York Avenue, New York, NY 10021 (US). GARIN-CHESA, Pilar; 1315 York Avenue, New York, NY 10021 (US). RETTIG, Wolfgang, J.; 1275 York Avenue, New York, NY 10021 (US). VAN DER BRUGGEN, Pierre; Avenue Hippocrate 74, UCL 7459, B- 1200 Brussels (BE). BOON-FALLEUR, Thierry; Avenue Hippocrate 74, UCL 7459, B-1200 Brussels (BE). OLD, Lloyd, J.; 1345 Avenue of the Americas, New York, NY 10105 (US).</p>		<p>(74) Agent: HANSON, Norman, D.; Felfe & Lynch, 805 Third Avenue, New York, NY 10022 (US). (81) Designated States: AU, CA, CN, FI, JP, NO, NZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>With amended claims.</i></p>
<p>(54) Title: MONOCLONAL ANBIBODIES WHICH BIND TO TUMOR REJECTION ANTIGEN PRECURSOR MAGE-1, RECOMBI- NANT MAGE-1, AND MAGE-1 DERIVED IMMUNOGENIC PEPTIDES (57) Abstract The invention relates to monoclonal antibodies which specifically bind to the tumor rejection antigen precursor molecule MAGE-1, hybridomas which produce these monoclonal antibodies, and their use. Also described is a recombinant form of MAGE-1, peptides which are useful as immunogens, and immunogenic compositions containing the peptides and an adjuvant.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5 MONOCLONAL ANTIBODIES WHICH BIND TO TUMOR REJECTION
ANTIGEN PRECURSOR MAGE-1, RECOMBINANT MAGE -1,
AND MAGE-1 DERIVED IMMUNOGENIC PEPTIDES

RELATED APPLICATION

10 This application is a continuation-in-part of Serial No.
037,230 filed March 26, 1993, which is itself a continuation-
in-part of PCT Application PCT/US92/04354 filed on May 22,
1992 designating the United States, which is a continuation-
in-part of Serial Number 807,043, filed December 12, 1991,
15 which is a continuation-in-part of Serial Number 764,365,
filed September 23, 1991, which is a continuation-in-part of
Serial Number 728,838, filed July 9, 1991, which is a
continuation-in-part of Serial Number 705,702, filed May 23,
1991, and now abandoned.

FIELD OF THE INVENTION

20 This invention relates in general to the field of immuno-
genetics as applied to the study of oncology. More
specifically, it relates to the study and analysis of
mechanisms by which tumors are recognized by the organism's
immune system such as through the presentation of so-called
25 tumor rejection antigens, and the expression of what will be
referred to herein as "tumor rejection antigen precursors" or
"TRAPs". Most specifically, it refers to one such TRAP, i.e.,
MAGE-1, produced recombinantly, and monoclonal antibodies and
antisera directed against MAGE-1, as well as their use.

30 BACKGROUND AND PRIOR ART

The study of the recognition or lack of recognition of
cancer cells by a host organism has proceeded in many
different directions. Understanding of the field presumes
some understanding of both basic immunology and oncology.

35 Early research on mouse tumors revealed that these
displayed molecules which led to rejection of tumor cells when
transplanted into syngeneic animals. These molecules are
"recognized" by T-cells in the recipient animal, and provoke
a cytolytic T-cell response with lysis of the transplanted
40 cells. This evidence was first obtained with tumors induced
in vitro by chemical carcinogens, such as methylcholanthrene.

5 The antigens expressed by the tumors and which elicited the T-cell response were found to be different for each tumor. See Prehn, et al., J. Natl. Canc. Inst. 18: 769-778 (1957); Klein et al., Cancer Res. 20: 1561-1572 (1960); Gross, Cancer Res. 3: 326-333 (1943), Basombrio, Cancer Res. 30: 2458-2462 (1970)
10 for general teachings on inducing tumors with chemical carcinogens and differences in cell surface antigens. This class of antigens has come to be known as "tumor specific transplantation antigens" or "TSTAs". Following the observation of the presentation of such antigens when induced
15 by chemical carcinogens, similar results were obtained when tumors were induced in vitro via ultraviolet radiation. See Kripke, J. Natl. Canc. Inst. 53: 333-1336 (1974).

While T-cell mediated immune responses were observed for the types of tumor described supra, spontaneous tumors were
20 thought to be generally non-immunogenic. These were therefore believed not to present antigens which provoked a response to the tumor in the tumor carrying subject. See Hewitt, et al., Brit. J. Cancer 33: 241-259 (1976).

The family of tum⁻ antigen presenting cell lines are
25 immunogenic variants obtained by mutagenesis of mouse tumor cells or cell lines, as described by Boon et al., J. Exp. Med. 152: 1184-1193 (1980), the disclosure of which is incorporated by reference. To elaborate, tum⁻ antigens are obtained by mutating tumor cells which do not generate an immune response
30 in syngeneic mice and will form tumors (i.e., "tum⁺" cells). When these tum⁺ cells are mutagenized, they are rejected by syngeneic mice, and fail to form tumors (thus "tum⁻"). See Boon et al., Proc. Natl. Acad. Sci. USA 74: 272 (1977), the disclosure of which is incorporated by reference. Many tumor
35 types have been shown to exhibit this phenomenon. See, e.g., Frost et al., Cancer Res. 43: 125 (1983).

It appears that tum⁻ variants fail to form progressive tumors because they elicit an immune rejection process. The evidence in favor of this hypothesis includes the ability of
40 "tum⁻" variants of tumors, i.e., those which do not normally form tumors, to do so in mice with immune systems suppressed

5 by sublethal irradiation, Van Pel et al., Proc. Natl, Acad.
Sci. USA 76: 5282-5285 (1979); and the observation that
intraperitoneally injected tum cells of mastocytoma P815
multiply exponentially for 12-15 days, and then are eliminated
in only a few days in the midst of an influx of lymphocytes
and macrophages (Uyttenhove et al., J. Exp. Med. 152: 1175-
1183 (1980)). Further evidence includes the observation that
mice acquire an immune memory which permits them to resist
subsequent challenge to the same tum variant, even when
immunosuppressive amounts of radiation are administered with
the following challenge of cells (Boon et al., Proc. Natl,
Acad. Sci. USA 74: 272-275 (1977); Van Pel et al., supra;
Uyttenhove et al., supra). Later research found that when
spontaneous tumors were subjected to mutagenesis, immunogenic
variants were produced which did generate a response. Indeed,
these variants were able to elicit an immune protective
response against the original tumor. See Van Pel et al., J.
Exp. Med. 157: 1992-2001 (1983). Thus, it has been shown that
it is possible to elicit presentation of a so-called "tumor
rejection antigen" in a tumor which is a target for a
syngeneic rejection response. Similar results have been
obtained when foreign genes have been transfected into
spontaneous tumors. See Fearson et al., Cancer Res. 48: 2975-
1980 (1988) in this regard.

A class of antigens has been recognized which are
presented on the surface of tumor cells and are recognized by
cytotoxic T cells, leading to lysis. This class of antigens
will be referred to as "tumor rejection antigens" or "TRAs"
hereafter. TRAs may or may not elicit antibody responses.
The extent to which these antigens have been studied, has been
via cytolytic T cell characterization studies in vitro i.e.,
the study of the identification of the antigen by a particular
cytolytic T cell ("CTL" hereafter) subset. The subset
proliferates upon recognition of the presented tumor rejection
antigen, and the cells presenting the antigen are lysed.
Characterization studies have identified CTL clones which
specifically lyse cells expressing the antigens. Examples of

5 this work may be found in Levy et al., Adv. Cancer Res. 24: 1-
59 (1977); Boon et al., J. Exp. Med. 152: 1184-1193 (1980);
Brunner et al., J. Immunol. 124: 1627-1634 (1980); Maryanski
et al., Eur. J. Immunol. 124: 1627-1634 (1980); Maryanski et
al., Eur. J. Immunol. 12: 406-412 (1982); Palladino et al.,
10 Canc. Res. 47: 5074-5079 (1987). This type of analysis is
required for other types of antigens recognized by CTLs,
including major histocompatibility antigens, the male specific
H-Y antigens, and a class of antigens, referred to as "tum-"
antigens, and discussed herein.

15 A tumor exemplary of the subject matter described supra
is known as P815. See DePlaen et al., Proc. Natl. Acad. Sci.
USA 85: 2274-2278 (1988); Szikora et al., EMBO J 9: 1041-1050
(1990), and Sibille et al., J. Exp. Med. 172: 35-45 (1990),
the disclosures of which are incorporated by reference. The
20 P815 tumor is a mastocytoma, induced in a DBA/2 mouse with
methylcholanthrene and cultured as both an in vitro tumor and
a cell line. The P815 line has generated many tum⁺ variants
following mutagenesis, including variants referred to as P91A
(DePlaen, supra), 35B (Szikora, supra), and P198 (Sibille,
25 supra). In contrast to tumor rejection antigens - and this is
a key distinction - the tum⁺ antigens are only present after
the tumor cells are mutagenized. Tumor rejection antigens are
present on cells of a given tumor without mutagenesis. Hence,
with reference to the literature, a cell line can be tum⁺,
30 such as the line referred to as "P1", and can be provoked to
produce tum⁺ variants. Since the tum⁺ phenotype differs from
that of the parent cell line, one expects a difference in the
DNA of tum⁺ cell lines as compared to their tum⁺ parental
lines, and this difference can be exploited to locate the gene
35 of interest in tum⁺ cells. As a result, it was found that
genes of tum⁺ variants such as P91A, 35B and P198 differ from
their normal alleles by point mutations in the coding regions
of the gene. See Szikora and Sibille, supra, and Lurquin et
al., Cell 58: 293-303 (1989). This has proved not to be the
40 case with the TRAs of this invention. These papers also
demonstrated that peptides derived from the tum⁺ antigen are

5 presented by the L^d molecule for recognition by CTLs. P91A is presented by L^d, P35 by D^d and P198 by K^d.

Prior patent applications PCT/US92/04354, U.S. Serial No. 807,043; 764,364; 728,838 and 707,702, all of which are incorporated by reference, describe inventions involving, inter alia, genes and other nucleic acid molecules which code for various TRAPs, which are in turn processed to tumor rejection antigen, or "TRAs".

10 The genes are useful as a source for the isolated and purified tumor rejection antigen precursor and the TRA themselves, either of which can be used as an agent for treating the cancer for which the antigen is a "marker", as well as in various diagnostic and surveillance approaches to oncology, discussed infra. It is known, for example, that tum⁻ cells can be used to generate CTLs which lyse cells presenting different tum⁻ antigens as well as tum⁺ cells. See, e.g., Maryanski et al., Eur. J. Immunol 12: 401 (1982); and Van den Eynde et al., Modern Trends in Leukemia IX (June 1990), the disclosures of which are incorporated by reference. The tumor rejection antigen precursor may be expressed in cells transfected by the gene, and then used to generate an immune response against a tumor of interest.

15 In the parallel case of human neoplasms, it has been observed that autologous mixed lymphocyte-tumor cell cultures ("MLTC" hereafter) frequently generate responder lymphocytes which lyse autologous tumor cells and do not lyse natural killer targets, autologous EBV-transformed B cells, or autologous fibroblasts (see Anichini et al., Immunol. Today 8: 385-389 (1987)). This response has been particularly well studied for melanomas, and MLTC have been carried out either with peripheral blood cells or with tumor infiltrating lymphocytes. Examples of the literature in this area including Knuth et al., Proc. Natl. Acad. Sci. USA 86: 2804-2802 (1984); Mukherji et al., J. Exp. Med. 158: 240 (1983); Hérin et al., Int. J. Canc. 39: 390-396 (1987); Topalian et al., J. Clin. Oncol 6: 839-853 (1988). Stable cytotoxic T cell clones ("CTLs" hereafter) have been derived from MLTC

5 responder cells, and these clones are specific for the tumor
cells. See Mukherji et al., supra, Hérin et al., supra, Knuth
et al., supra. The antigens recognized on tumor cells by
these autologous CTLs do not appear to represent a cultural
10 artifact, since they are found on tumor cells in vivo.
Topalian et al., supra; Degiovanni et al., Eur. J. Immunol.
20: 1865-1868 (1990). These observations, coupled with the
techniques used herein to isolate the genes for specific
murine tumor rejection antigen precursors, have led to the
15 isolation of nucleic acid sequences coding for tumor rejection
antigen precursors of TRAs presented on human tumors. It is
now possible to isolate the nucleic acid sequences which code
for tumor rejection antigen precursors, including, but not
being limited to those most characteristic of a particular
tumor, with ramifications that are described infra.

20 Additional work has focused upon the presentation of TRAs
by the class of molecules known as human leukocyte antigens,
or "HLAs". This work has resulted in several unexpected
discoveries regarding the field. Specifically in U.S. patent
application Serial Number 938,334, the disclosure of which is
25 incorporated by reference, nonapeptides are taught which are
presented by the HLA-A1 molecule. The reference teaches that
given the known specificity of particular peptides for
particular HLA molecules, one should expect a particular
peptide to bind one HLA molecule, but not others. This is
30 important, because different individuals possess different HLA
phenotypes. As a result, while identification of a particular
peptide as being a partner for a specific HLA molecule has
diagnostic and therapeutic ramifications, these are only
relevant for individuals with that particular HLA phenotype.
35 There is a need for further work in the area, because cellular
abnormalities are not restricted to one particular HLA
phenotype, and targeted therapy requires some knowledge of the
phenotype of the abnormal cells at issue.

40 In U.S. Patent Application Serial Number 008,446, filed
January 22, 1993 and incorporated by reference, the fact that
the MAGE-1 expression product is processed to a second TRA is

5 disclosed. This second TRA is presented by HLA-C-clone-10 molecules. The disclosure shows that a given TRAP can yield a plurality of TRAs.

10 In U.S. Patent Application Serial Number 994,928, filed December 22, 1992, and incorporated by reference herein, tyrosinase is described as a tumor rejection antigen precursor. This reference discloses that a molecule which is produced by some normal cells (e.g., melanocytes), is processed in tumor cells to yield a tumor rejection antigen that is presented by HLA-A2 molecules.

15 The prior applications cited supra discussed antibodies against tumor rejection antigen precursors generally. The present investigators have utilized the isolated nucleic acid molecule coding for MAGE-1 to produce a recombinant MAGE-1 protein, and peptides derived therefrom. These have been used
20 to produce polyclonal and monoclonal antibodies which specifically bind to MAGE-1. These antibodies, and their use, constitute the invention described and claimed herein.

BRIEF DESCRIPTION OF THE FIGURES

25 Figure 1 shows, schematically, the MAGE-1 gene, oligopeptides derived from the recombinant MAGE-1 protein, and comparison with corresponding sequences in MAGE-2 and MAGE-3 deduced amino acid sequences.

30 Figure 2A shows silver stained SDS-polyacrylamide gels of affinity purified, MAGE-1 recombinant protein. Figure 2B presents immunoblotting work where recombinant MAGE-1 protein was used against rabbit antisera derived from immunization with three peptides (SEQ ID NOS: 2, 3 and 4). Blotting was at 1:1000 dilution. As a control, recombinant mouse p53 was used.

35 Figure 3A shows the reactivity pattern of mAb MA 454 against six melanoma lines. Figure 3B shows the results obtained using rabbit polyclonal antisera against the same lines. Figure 3C shows results obtained with a MAGE-1 transfected cell line (MZ2-MEL 2.2-ET.1), and its parent (MZ2-MEL 2.2).
40

Figure 4 presents immunoblot analysis using the

antibodies against tissue lysates.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Many different "MAGE" genes have been identified, as will be seen from the applications and references cited *supra*. The MAGE-1 gene is at issue on the present case, and is the only one discussed hereafter. For convenience, it is presented herein as SEQ ID NO: 1.

"MAGE" as used herein refers to a nucleic acid sequence isolated from human cells.

When "TRAP" or "TRAS" are discussed herein as being specific to a tumor type, this means that the molecule under consideration is associated with that type of tumor, although not necessarily to the exclusion of other tumor types.

Example 1

The cell line MZ2-MEL 3.1 described in, e.g., Van den Eynde et al., Int. J. Cancer 44: 634-640 (1989) and in the parent application cited *supra*, previously observed to express MAGE-1, was used as a source of total RNA. The total RNA was extracted from the cells, and was then subjected to reverse transcription/polymerase chain reaction, using the primers CH08 and CH09, as described by Van der Bruggen et al., Science 254: 1643-1647 (Dec. 13, 1991), the disclosure of which is incorporated by reference. This paper describes the "RT-PCR" technique, as does the Brasseur et al., Int. J. Cancer 52: 839-841 (1992). It must be understood, however, that the sequence of MAGE-1 is known to the art, and other primers could be used besides CH08 and CH09.

Once the RT-PCR protocols were completed, the products were cloned directly into plasmid pT7 Blue (Novagen, Madison WI), following manufacturer's instructions which constituted well known techniques. Following the cloning, the recombinant plasmid DNA was treated with restriction endonucleases to generate fragments which included fragments containing the MAGE-1 gene. See, e.g. Van der Bruggen et al, *supra*.

The appropriate cDNA insert was subcloned unidirectionally, into plasmids pQE9, pQE10 and pQE11, using BamHI and HindIII cloning sites in pT7 Blue. The plasmids

5 were transfected into *E. coli*, and recombinant protein production was induced via IPTG, as the host plasmid contains the lac operon. This yielded a fusion protein containing the MAGE-1 polypeptide sequence, which could be purified via Ni²⁺ ion affinity chromatography.

10 The DNA sequence of the recombinant clone was obtained, and was confirmed to encode 163 amino acids which correspond to deduced amino acids 57-219 of predicted MAGE-1 amino sequence, plus 30 residues from the plasmid itself. Figure 1 shows this. The expected molecular mass is about 20-22 kDa.

15 When clones in pQE10 were studied, indeed, a recombinant protein of about 20 kDa was produced following IPTG induction. Other minor protein species of 70 kDa, 43 kDa, 17 kDa and 15 kDa were also found, as is seen in figure 2A.

Example 2

20 The following describes procedures used to produce antibodies to MAGE-1. Based upon the predicted MAGE-1 amino acid sequence, three oligopeptides were prepared:

25 Ile Asn Phe Thr Arg Gln Arg Gln Pro Ser Glu Gly Ser Ser
(SEQ ID NO: 2)

Leu Phe Arg Ala Val Ile Thr Lys Lys Val Ala Asp
(SEQ ID NO: 3)

30 Asp Val Lys Glu Ala Asp Pro Thr Gly His Ser Tyr
(SEQ ID NO: 4)

Rabbits were immunized with the peptides, and were then treated to collect antiserum.

35 Antisera prepared against these three peptides were then used with *E. coli* produced, recombinant MAGE-1 protein, in immunoblotting experiments. The results, set forth in figure 2B, show that only antiserum raised against the first of these peptides, i.e., SEQ ID NO: 2 reacted strongly. The fact that
40 additional protein species that copurified with the 20 kDa fusion protein also showed reactivity, suggests that these are

5 aggregates of the fusion protein. The peptide used corresponds to deduced amino acids 68-81 of the MAGE-1 of the predicted MAGE-1 protein.

10 When immunoblotting was carried out using lysates of melanoma cell line MZ2-MEL 3.1, no detectable MAGE protein was found.

Example 3

15 Monoclonal antibodies were then prepared. Purified recombinant protein, produced as described supra, was used to immunize BALB/C mice. Hybridomas were generated and cloned. The protocol used was that described by Dippold et al., Proc. Natl. Acad. Sci. USA 77: 6114-6118 (1980), the disclosure of which is incorporated by reference. The key difference, of course, was the immunogen used for immunization.

20 Once hybridomas were generated, their supernatants were screened using a standard, solid phase ELISA on microtiter plates, using the immunizing fusion protein as target antigen. Five clones were found to be reactive. They all also showed moderate to strong reactivity in immunoblots.

25 As a control, mouse p53 protein, expressed in the same plasmid vector, was also tested. No reactivity was seen. These results are summarized in Table 1 which follows:

5 TABLE 1. Reactivity of mouse anti-recombinant MAGE-1, mAbs
toward recombinant MAGE-1 protein and control p53 protein

Assay mAb	ELISA		Immunoblot	
	MAGE-1	p53	MAGE-1	p53
MA32	+++	-	++ [#]	-
MA231	+	-	++	-
MA399	++	-	++	-
MA430	++	-	+++	-
MA454	++	-	+++	-

*ELISA titer using hybridoma supernatants: -, <1:16; +, 1:64; ++, 1:256. #Immunoblot signal intensity: -, negative; +, weak; ++, moderate; +++, strong.

Example 4

The mAbs described supra were then tested against lysates of melanoma cell lines. The cell lines tested, i.e., MZ2-MEL 3.1, MZ2-MEL 2.2, and SK-MEL-187, are all well known. MZ2-MEL 2.2 is a MAGE-1 loss variant derived from MAGE-1 positive parental MZ2-MEL 3.1 by CTL selection (van der Bruggen et al., Int. J. Cancer 44: 634-640 (1989)). These cells had been "typed" by RT-PCR as being MAGE-1⁺2⁺3⁺ (MZ2-MEL 3.1), MAGE-1⁺2⁺3⁻ (MZ2-MEL 2.2), and MAGE 1⁻2⁻3⁻ (SK MEL-187). The lysates were prepared by homogenizing the cells in Nonidet P40 (NP-40) buffer (1% NP-40, 50 mM Tris-HCl, pH 8.0, 150 mM NaCl). The results are shown in figure 3A.

Monoclonal antibody MA 454 reacted with a 46 kDa protein present in MZ2-MEL 3.1 lysate, but not in lysates of either of the other two cell lines. When three additional melanoma

5 lines were tested, only those which were typed as being MAGE-1 positive reacted with the mAb. Expression of MAGE-2 or MAGE-3 was irrelevant.

10 The polyclonal antiserum described supra, was also tested against these lysates. Results are shown in figure 3B. It was positive for MZ2-MEL 3.1, and for MAGE-1 transfected cell line MZ2-MEL 2.2-ET.1, but was negative for parental line MZ2-MEL 2.2.

Example 5

15 Lysates were prepared from liver, kidney and testis tissue, and from four melanoma cell lines including one MAGE-1²3³ line, two MAGE-1²3³, and one MAGE-1²3³ lines. The lysates were prepared as described supra. When immunoblotting was carried out, testis lysates were positive with mAb 454, as were MAGE-1 positive melanomas. No other lysates were
20 positive which is in complete agreement with mRNA expression data.

The same experiments were carried out using polyclonal antiserum, and the results paralleled those for the monoclonal antibodies. Figure 4 presents these results.

25 The foregoing experiments describe the production of monoclonal antibodies which specifically bind to a tumor rejection antigen precursor TRAP. The studies show binding both to the "wild type" MAGE-1 molecule, and the recombinant form, but not to either of MAGE-2 or MAGE-3. A particularly preferred species of MAGE-1 binding mAb, i.e., MA454, has been
30 deposited at the American Type Culture Collection under Accession Number HB 11540.

35 The invention thus relates to MAGE-1 specific monoclonal antibodies and the hybridomas which produce them. The mAbs were found to be useful in determining expression of MAGE-1 in cell lysates. Specifically, the mAbs can be added, e.g., in labelled form, bound to a solid phase, or otherwise treated to increase the sensitivity of MAGE-1 detection. Any of the standard types of immunoassays, including ELISAs, RIAs, competitive assays, agglutination assays, and all others are
40 encompassed with respect to the way the mAbs can be used.

5 "Cell lysate" as used herein refers not only to a sample which
is expressly lysed, but also to those samples which contain
cells which have been lysed in vivo, or any sample which
contains material normally internal to the cells. The
detection of MAGE-1 expression product is useful, e.g., in
10 diagnosing or monitoring the presence or progress of a cancer.

The isolated, recombinant MAGE-1 protein is also a
feature of this invention. This molecule has a molecular
weight of about 20-22 kDa as determined by SDS-PAGE, and is
useful as an immunogen as are the peptides of SEQ ID NOS: 2,
15 3 and 4, shown by the examples to be immunogenic.
Preferably, these are used in combination with a suitable
adjuvant. The isolated form of the molecule obtained via non-
recombinant means has a molecular weight of about 43 kd as
determined by SDS-PAGE, and is useful in the same ways as is
20 the recombinant protein. The recombinant form may consist of
only amino acids 57-219 of the sequence of MAGE-1, as shown
supra. Also a part of the invention is the full length
isolated, recombinant MAGE-1 protein, having a molecular
weight of about 34.3kd as determined by SDS-PAGE, and
25 consisting of the amino acid sequence coded for by nucleotides
3931-4761 of SEQ ID NO: 1.

Other features of the invention will be clear to the
artisan and need not be repeated here.

30 The terms and expressions which have been employed are
used as terms of description and not of limitation, and there
is no intention in the use of such terms and expressions of
excluding any equivalents of the features shown and described
or portions thereof, it being recognized that various
modifications are possible within the scope of the invention.
35

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANTS: Chen, Yao-Tseng; Stockert, Elisabeth;
Chen, Yachi; Garin-Chesa, Pilar; Rettig, Wolfgang
J.; van der Bruggen, Pierre; Boon-Falleur,
Thierry; Old, Lloyd J.

(ii) TITLE OF INVENTION: MONOCLONAL ANTIBODIES WHICH
BIND TO TUMOR REJECTION ANTIGEN PRECURSOR MAGE-1,
RECOMBINANT MAGE-1, AND MAGE-1 DERIVED IMMUNOGENIC
PEPTIDES

(iii) NUMBER OF SEQUENCES: 4

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Felfe & Lynch
(B) STREET: 805 Third Avenue
(C) CITY: New York City
(D) STATE: New York
(F) ZIP: 10022

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette, 5.25 inch, 360 kb
storage

(B) COMPUTER: IBM
(C) OPERATING SYSTEM: PC-DOS
(D) SOFTWARE: Wordperfect

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: 08/190,411
(B) FILING DATE: 01-FEBRUARY-1994
(C) CLASSIFICATION: 424

(vii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: 037,230
- (B) FILING DATE: 26-MARCH-1993

(vii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: PCT/US92/04354
- (B) FILING DATE: 22-MAY-1992

(viii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: 07/807,043
- (B) FILING DATE: 12-DECEMBER-1991

(ix) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: 07/764,364
- (B) FILING DATE: 23-SEPTEMBER-1991

(x) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: 07/728,838
- (b) FILING DATE: 9-JULY-1991

(xi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: 07/705,702
- (B) FILING DATE: 23-MAY-1991

(xii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Hanson, Norman D.
- (B) REGISTRATION NUMBER: 30,946
- (C) REFERENCE/DOCKET NUMBER: LUD 5354

(xiii) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (212) 688-9200
- (B) TELEFAX: (212) 838-3884

(2) INFORMATION FOR SEQUENCE ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5674 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: genomic DNA

(ix) FEATURE:

(A) NAME/KEY: MAGE-1 gene

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CCCGGGGCAC	CACTGGCATC	CCTCCCCCTA	CCACCCCCAA	TCCCTCCCTT	50
TACGCCACCC	ATCCAAACAT	CTTCACGCTC	ACCCCCAGCC	CAAGCCAGGC	100
AGAATCCGGT	TCCACCCCTG	CTCTCAACCC	AGGGAAGCCC	AGGTGCCCAG	150
ATGTGACGCC	ACTGACTTGA	GCATTAGTGG	TTAGAGAGAA	GCGAGGTTTT	200
CGGTCTGAGG	GGCGGCTTGA	GATCGGTGGA	GGGAAGCGGG	CCCAGCTCTG	250
TAAGGAGGCA	AGGTGACATG	CTGAGGGAGG	ACTGAGGACC	CACTTACCCC	300
AGATAGAGGA	CCCCAAATAA	TCCCTTCATG	CCAGTCCTGG	ACCATCTGGT	350
GGTGGACTTC	TCAGGCTGGG	CCACCCCCAG	CCCCCTTGCT	GCTTAAACCA	400
CTGGGGACTC	GAAGTCAGAG	CTCCGTGTGA	TCAGGGAAGG	GCTGCTTAGG	450
AGAGGGCAGC	GTCCAGGCTC	TGCCAGACAT	CATGCTCAGG	ATTCTCAAGG	500
AGGGCTGAGG	GTCCCTAAGA	CCCCACTCCC	GTGACCCAAC	CCCCACTCCA	550
ATGCTCACTC	CCGTGACCCA	ACCCCCTCTT	CATTGTCATT	CCAACCCCCA	600
CCCCACATCC	CCCACCCCAT	CCCTCAACCC	TGATGCCCAT	CCGCCCAGCC	650
ATTCCACCCT	CACCCCCACC	CCCACCCCCA	CGCCCCACTC	CACCCCCACC	700
CAGGCAGGAT	CCGGTTCCCG	CCAGGAAACA	TCCGGGTGCC	CGGATGTGAC	750
GCCACTGACT	TGCGCATTGT	GGGGCAGAGA	GAAGCGAGGT	TTCCATTCTG	800
AGGGACGGCG	TAGAGTTCGG	CCGAAGGAAC	CTGACCCAGG	CTCTGTGAGG	850
AGGCAAGGTG	AGAGGCTGAG	GGAGGACTGA	GGACCCCGCC	ACTCCAAATA	900
GAGAGCCCCA	AATATTCCAG	CCCCGCCCTT	GCTGCCAGCC	CTGGCCCACC	950
CGCGGGAAGA	CGTCTCAGCC	TGGGCTGCCC	CCAGACCCCT	GCTCCAAAAG	1000
CCTTGAGAGA	CACCAGGTTC	TTCTCCCCAA	GCTCTGGAAT	CAGAGGTTGC	1050
TGTGACCAGG	GCAGGACTGG	TTAGGAGAGG	GCAGGGCACA	GGCTCTGCCA	1100
GGCATCAAGA	TCAGCACCCA	AGAGGGAGGG	CTGTGGGCCC	CCAAGACTGC	1150
ACTCCAATCC	CCACTCCCAC	CCCATTGCGA	TTCCCATTCC	CCACCCAACC	1200
CCCATCTCCT	CAGCTACACC	TCCACCCCCA	TCCCTACTCC	TACTCCGTCA	1250

CCTGACCACC	ACCCTCCAGC	CCCAGCACCA	GCCCCAACCC	TTCTGCCACC	1300
TCACCCTCAC	TGCCCCCAAC	CCCACCCTCA	TCTCTCTCAT	GTGCCCCACT	1350
CCCATCGCCT	CCCCCATTTCT	GGCAGAATCC	GGTTTGCCCC	TGCTCTCAAC	1400
CCAGGGAAGC	CCTGGTAGGC	CCGATGTGAA	ACCACTGACT	TGAACCTCAC	1450
AGATCTGAGA	GAAGCCAGGT	TCATTTAATG	GTTCTGAGGG	GCGGCTTGAG	1500
ATCCACTGAG	GGGAGTGGTT	TTAGGCTCTG	TGAGGAGGCA	AGGTGAGATG	1550
CTGAGGGAGG	ACTGAGGAGG	CACACACCCC	AGGTAGATGG	CCCCAAAATG	1600
ATCCAGTACC	ACCCTTGCTG	CCAGCCCTGG	ACCACCCGGC	CAGGACAGAT	1650
GTCTCAGCTG	GACCACCCCC	CGTCCCGTCC	CACTGCCACT	TAACCCACAG	1700
GGCAATCTGT	AGTCATAGCT	TATGTGACCG	GGGCAGGGTT	GGTCAGGAGA	1750
GGCAGGGCCC	AGGCATCAAG	GTCCAGCATC	CGCCCGGCAT	TAGGGTCAGG	1800
ACCCTGGGAG	GGAAGTGAAG	GTTCCCCACC	CACACCTGTC	TCCTCATCTC	1850
CACCGCCACC	CCACTCACAT	TCCCATACCT	ACCCCTACC	CCCAACCTCA	1900
TCTTGTCAGA	ATCCCTGCTG	TCAACCCACG	GAAGCCACGG	GAATGGCGGC	1950
CAGGCACTCG	GATCTTGACG	TCCCCATCCA	GGGTCTGATG	GAGGGAAGGG	2000
GCTTGAACAG	GGCCTCAGGG	GAGCAGAGGG	AGGGCCCTAC	TGCGAGATGA	2050
GGGAGGCCTC	AGAGGACCCA	GCACCCTAGG	ACACCGCACC	CCTGTCTGAG	2100
ACTGAGGCTG	CCACTTCTGG	CCTCAAGAAT	CAGAACGATG	GGGACTCAGA	2150
TTGCATGGGG	GTGGGACCCA	GGCCTGCAAG	GCTTACGCGG	AGGAAGAGGA	2200
GGGAGGACTC	AGGGGACCTT	GGAATCCAGA	TCAGTGTGGA	CCTCGGCCCT	2250
GAGAGGTCCA	GGGCACGGTG	GCCACATATG	GCCCATATTT	CCTGCATCTT	2300
TGAGGTGACA	GGACAGAGCT	GTGGTCTGAG	AAGTGGGGCC	TCAGGTCAAC	2350
AGAGGGAGGA	GTTCCAGGAT	CCATATGGCC	CAAGATGTGC	CCCCTTCATG	2400
AGGACTGGGG	ATATCCCCGG	CTCAGAAAGA	AGGGACTCCA	CACAGTCTGG	2450
CTGTCCCCTT	TTAGTAGCTC	TAGGGGGACC	AGATCAGGGA	TGGCGGTATG	2500
TTCCATTCTC	ACTTGTAACA	CAGGCAGGAA	GTTGGGGGGC	CCTCAGGGAG	2550
ATGGGGTCTT	GGGGTAAAGG	GGGGATGTCT	ACTCATGTCA	GGGAATTGGG	2600
GGTTGAGGAA	GCACAGGCGC	TGGCAGGAAT	AAAGATGAGT	GAGACAGACA	2650
AGGCTATTGG	AATCCACACC	CCAGAACCA	AGGGGTCAGC	CCTGGACACC	2700
TCACCCAGGA	TGTGGCTTCT	TTTTCACTCC	TGTTTCCAGA	TCTGGGGCAG	2750
GTGAGGACCT	CATTCTCAGA	GGGTGACTCA	GGTCAACGTA	GGGACCCCCA	2800
TCTGGTCTAA	AGACAGAGCG	GTCCCAGGAT	CTGCCATGCG	TTCGGGTGAG	2850
GAACATGAGG	GAGGACTGAG	GGTACCCAG	GACCAGAACA	CTGAGGGAGA	2900
CTGCACAGAA	ATCAGCCCTG	CCCCTGCTGT	CACCCAGAG	AGCATGGGCT	2950
GGGCCGTCTG	CCGAGGTCCT	TCCGTTATCC	TGGGATCATT	GATGTCAGGG	3000
ACGGGGAGGC	CTTGGTCTGA	GAAGGCTGCG	CTCAGGTCAG	TAGAGGGAGC	3050
GTCCCAGGCC	CTGCCAGGAG	TCAAGGTGAG	GACCAAGCGG	GCACCTCACC	3100

CAGGACACAT² TAATTCCAAT GAATTTTGAT ATCTCTTGCT GCCCTTCCCC 3150
 AAGGACCTAG GCACGTGTGG CCAGATGTTT GTCCCCCTCCT GTCCTTCCAT 3200
 TCCTTATCAT GGATGTGAAC TCTTGATTTG GATTTCTCAG ACCAGCAAAA 3250
 GGGCAGGATC CAGGCCCTGC CAGGAAAAAT ATAAGGGCCC TGC GTGAGAA 3300
 CAGAGGGGGT CATCCACTGC ATGAGAGTGG GGATGTCACA GAGTCCAGCC 3350
 CACCCTCCTG GTAGCACTGA GAAGCCAGGG CTGTGCTTGC GGTCTGCACC 3400
 CTGAGGGGCC GTGGATTCTT CTTCCTGGAG CTCCAGGAAC CAGGCAGTGA 3450
 GGCCTTGGTC TGAGACAGTA TCCTCAGGTC ACAGAGCAGA GGATGCACAG 3500
 GGTGTGCCAG CAGTGAATGT TTGCCCTGAA TGCACACCAA GGGCCCCACC 3550
 TGCCACAGGA CACATAGGAC TCCACAGAGT CTGGCCTCAC CTCCCTACTG 3600
 TCAGTCCTGT AGAATCGACC TCTGCTGGCC GGCTGTACCC TGAGTACCCT 3650
 CTCACTTCCT CCTTCAGGTT TTCAGGGGAC AGGCCAACCC AGAGGACAGG 3700
 ATTCCCTGGA GGCCACAGAG GAGCACCAAG GAGAAGATCT GTAAGTAGGC 3750
 CTTTGTTAGA GTCTCCAAGG TTCAGTTCTC AGCTGAGGCC TCTCACACAC 3800
 TCCCTCTCTC CCCAGGCCTG TGGGTCTTCA TTGCCCAGCT CCTGCCCACA 3850
 CTCCTGCCTG CTGCCCTGAC GAGAGTCATC 3880
 ATG TCT CTT GAG CAG AGG AGT CTG CAC TGC AAG CCT GAG GAA 3922
 GCC CTT GAG GCC CAA CAA GAG GCC CTG GGC CTG GTG TGT GTG 3964
 CAG GCT GCC ACC TCC TCC TCC TCT CCT CTG GTC CTG GGC ACC 4006
 CTG GAG GAG GTG CCC ACT GCT GGG TCA ACA GAT CCT CCC CAG 4048
 AGT CCT CAG GGA GCC TCC GCC TTT CCC ACT ACC ATC AAC TTC 4090
 ACT CGA CAG AGG CAA CCC AGT GAG GGT TCC AGC AGC CGT GAA 4132
 GAG GAG GGG CCA AGC ACC TCT TGT ATC CTG GAG TCC TTG TTC 4174
 CGA GCA GTA ATC ACT AAG AAG GTG GCT GAT TTG GTT GGT TTT 4216
 CTG CTC CTC AAA TAT CGA GCC AGG GAG CCA GTC ACA AAG GCA 4258
 GAA ATG CTG GAG AGT GTC ATC AAA AAT TAC AAG CAC TGT TTT 4300
 CCT GAG ATC TTC GGC AAA GCC TCT GAG TCC TTG CAG CTG GTC 4342
 TTT GGC ATT GAC GTG AAG GAA GCA GAC CCC ACC GGC CAC TCC 4384
 TAT GTC CTT GTC ACC TGC CTA GGT CTC TCC TAT GAT GGC CTG 4426
 CTG GGT GAT AAT CAG ATC ATG CCC AAG ACA GGC TTC CTG ATA 4468
 ATT GTC CTG GTC ATG ATT GCA ATG GAG GGC GGC CAT GCT CCT 4510
 GAG GAG GAA ATC TGG GAG GAG CTG AGT GTG ATG GAG GTG TAT 4552
 GAT GGG AGG GAG CAC AGT GCC TAT GGG GAG CCC AGG AAG CTG 4594
 CTC ACC CAA GAT TTG GTG CAG GAA AAG TAC CTG GAG TAC GGC 4636
 AGG TGC CGG ACA GTG ATC CCG CAC GCT ATG AGT TCC TGT GGG 4678
 GTC CAA GGG CCC TCG CTG AAA CCA GCT ATG TGA 4711
 AAGTCCTTGA GTATGTGATC AAGGTCAGTG CAAGAGTTC 4750

GCTTTTTCTT	CCCATCCCTG	CGTGAAGCAG	CTTTGAGAGA	GGAGGAAGAG	4800
GGAGTCTGAG	CATGAGTTGC	AGCCAAGGCC	AGTGGGAGGG	GGACTGGGCC	4850
AGTGCACCTT	CCAGGGCCGC	GTCCAGCAGC	TTCCCCTGCC	TCGTGTGACA	4900
TGAGGCCCAT	TCTTCACTCT	GAAGAGAGCG	GTCAGTGTTT	TCAGTAGTAG	4950
GTTTCTGTTC	TATTGGGTGA	CTTGGAGATT	TATCTTTGTT	CTCTTTTGGA	5000
ATTGTTCAAA	TGTTTTTTTT	TAAGGGATGG	TTGAATGAAC	TTCAGCATCC	5050
AAGTTTATGA	ATGACAGCAG	TCACACAGTT	CTGTGTATAT	AGTTTAAGGG	5100
TAAGAGTCTT	GTGTTTTATT	CAGATTGGGA	AATCCATTCT	ATTTTGTGAA	5150
TTGGGATAAT	AACAGCAGTG	GAATAAGTAC	TTAGAAATGT	GAAAAATGAG	5200
CAGTAAAATA	GATGAGATAA	AGAACTAAAG	AAATTAAGAG	ATAGTCAATT	5250
CTTGCCTTAT	ACCTCAGTCT	ATTCTGTAAA	ATTTTTAAAG	ATATATGCAT	5300
ACCTGGATTT	CCTTGGCTTC	TTTGAGAATG	TAAGAGAAAT	TAAATCTGAA	5350
TAAAGAATTC	TTCCTGTTCA	CTGGCTCTTT	TCTTCTCCAT	GCACTGAGCA	5400
TCTGCTTTTT	GGAAGGCCCT	GGGTTAGTAG	TGGAGATGCT	AAGGTAAGCC	5450
AGACTCATAC	CCACCCATAG	GGTCGTAGAG	TCTAGGAGCT	GCAGTCACGT	5500
AATCGAGGTG	GCAAGATGTC	CTCTAAAGAT	GTAGGGAAAA	GTGAGAGAGG	5550
GGTGAGGGTG	TGGGGCTCCG	GGTGAGAGTG	GTGGAGTGTC	AATGCCCTGA	5600
GCTGGGGCAT	TTTGGGCTTT	GGGAAACTGC	AGTTCCTTCT	GGGGGAGCTG	5650
GCTGGGGCAT	TTTGGGCTTT	GGGAAACTGC	AGTTCCTTCT	GGGGGAGCTG	5700
ATTGTAATGA	TCTTGGGTGG	ATCC			5724

20

(2) INFORMATION FOR SEQUENCE ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acid residues
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Ile Asn Phe Thr Arg Gln Arg Gln Pro Ser Glu Gly Ser Ser
5 10

(2) INFORMATION FOR SEQUENCE ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acid residues
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Leu Phe Arg Ala Val Ile Thr Lys Lys Val Ala Asp
5 10

(2) INFORMATION FOR SEQUENCE ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acid residues
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Asp Val Lys Glu Ala Asp Pro Thr Gly His Ser Tyr
5 10

We claim:

1. Monoclonal antibody which specifically binds to tumor rejection antigen precursor MAGE-1.
2. The monoclonal antibody of claim 1, designated MA454.
3. Hybridoma cell line which produces the monoclonal antibody of claim 1.
4. The hybridoma cell line of claim 3, wherein said monoclonal antibody is MA454.
5. Method for determining tumor rejection antigen precursor MAGE-1 in a sample, comprising contacting said sample with the monoclonal antibody of claim 1 and determining binding of said monoclonal antibody to a component of said sample as a determination of MAGE-1 in said sample.
6. The method of claim 5, wherein said monoclonal antibody is bound to a solid phase.
7. The method of claim 5, wherein said monoclonal antibody is labelled with a detectable label.
8. Isolated, MAGE-1 tumor rejection antigen precursor.
9. The isolated MAGE-1 tumor rejection antigen precursor of claim 8, which is a glycoprotein having a molecular weight of about 46 kilodaltons as determined by SDS-PAGE.
10. The isolated MAGE-1 tumor rejection antigen precursor of claim 8, which is a recombinantly produced protein having a molecular weight of about 34.3 kilodaltons as determined by SDS-PAGE.

11. Isolated protein consisting of amino acids 57-219 coded for by nucleotides 3931-4761 of the nucleotide sequence of SEQ I.D. NO.: 1.
12. Isolated peptide selected from the group consisting of:
SEQ ID NO: 2;
SEQ ID NO: 3, and
SEQ ID NO: 4.
13. Immunogenic composition comprising at least one isolated protein of claim 9 and an adjuvant.
14. Immunogenic composition comprising at least one isolated protein of claim 10 and an adjuvant.
15. Immunogenic composition comprising at least one isolated protein of claim 11 and an adjuvant.
16. Immunogenic composition comprising at least one peptide of claim 12 and an adjuvant.

AMENDED CLAIMS

[received by the International Bureau on 9 May 1995 (09.05.95);
original claims 1-7 unchanged; original claims 8-16 replaced by
new claims 8-13 (2 pages)]

1. Monoclonal antibody which specifically binds to tumor rejection antigen precursor MAGE-1.
2. The monoclonal antibody of claim 1, designated MA454.
3. Hybridoma cell line which produces the monoclonal antibody of claim 1.
4. The hybridoma cell line of claim 3, wherein said monoclonal antibody is MA454.
5. Method for determining tumor rejection antigen precursor MAGE-1 in a sample, comprising contacting said sample with the monoclonal antibody of claim 1 and determining binding of said monoclonal antibody to a component of said sample as a determination of MAGE-1 in said sample.
6. The method of claim 5, wherein said monoclonal antibody is bound to a solid phase.
7. The method of claim 5, wherein said monoclonal antibody is labelled with a detectable label.
8. The isolated MAGE-1 tumor rejection antigen precursor derivative which is a protein having a molecular mass of from about 20 kilodaltons to about 22 kilodaltons.
9. Isolated protein consisting of amino acids 57-219 coded for by nucleotides 3931-4761 of the nucleotide sequence of SEQ ID NO: 1.

AMENDED SHEET (ARTICLE 19)


10. Isolated peptide selected from the group consisting of:

SEQ ID NO: 2,
SEQ ID NO: 3, and
SEQ ID NO: 4.

11. Immunogenic composition comprising at least one isolated protein of claim 8 and an adjuvant.
12. Immunogenic composition comprising at least one isolated protein of claim 9 and an adjuvant.
13. Immunogenic composition comprising at least one isolated peptide of claim 10 and an adjuvant.

AMENDED SHEET (ARTICLE 19)



Fig. 1




	pep I	pep II	pep III
MAGE-1	INFTRQRPSEGS	LFRAVITKKVAD	DVKEADPTGHSY
MAGE-2	--Y-LW--SD---	E-Q-A-SR-MVE	E-V-VV-I-S-L-
MAGE-3	M-YPLW S SY-D-	E-Q-ALSR--E	ELM-V--I--L-

SUBSTITUTE SHEET (RULE 26)

2/5


Fig. 2B*Fig. 2A*

SUBSTITUTE SHEET (RULE 26)



SUBSTITUTE SHEET (RULE 26)

5/5

*Fig. 46*Rabbit anti-MAGE-1

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/00095

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/ 277.1; 435/7.1, 7.23, 240.27; 530/324, 327, 350, 387.7, 387.9, 388.8, 389.7, 391.1, 391.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CA, MEDLINE, INPADOC, EMBASE, BIOSIS, REGISTRY, HCAPLUS

search terms: MAGE, antigen, peptide, protein, polypeptide, antigen, INFTRQRPSEGSS, LFRAVITKKVAD, DVKEADPTGHS, antibod?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	WO, A, 92/20356 (BOON ET AL) 26 November 1992, see entire document.	8 ----- 1-7, 9-11
P, X	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, Volume 91, issued February 1994, Y. Chen et al., "Identification of the MAGE-1 Gene Product by Monoclonal and Polyclonal Antibodies", pages 1004-1008, see entire document.	1-5, 8-11
P, Y	MOLECULAR IMMUNOLOGY, Volume 31, Number 18, issued 1994. E. Celis et al., "Identification of Potential CTL Epitopes of Tumor Associated Antigen MAGE-1 for Five Common HLA-Alleles", pages 1423-1430, see entire document.	12-16

☒ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z

document member of the same patent family

Date of the actual completion of the international search

27 FEBRUARY 1995

Date of mailing of the international search report

03 APR 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ANTHONY C. CAPUTA, Ph.D.

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/00095

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	JOURNAL OF EXPERIMENTAL MEDICINE. Volume 176, issued November 1992, C. Traversari et al., "A Nonapeptide Encoded by Human Gene MAGE-1 is Recognized on HLA-A1 by Cytolytic T Lymphocytes Directed Against Tumor Antigen MZ2-E", pages 1453-1457, see entire document.	12-16
Y	E. HARLOW et al, ANTIBODIES, A LABORATORY MANUAL", published 1988 by Cold Spring Harbor Laboratory (N.Y.), pages 96, 97, 139, 140, 148, 149, 553-556, and 578-582, see pages 96, 97, 139, 140, 148, 149, 553-556, and 578-582.	1-7, 13-16

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No
PCT/US95/00095

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (6):

A61K 38/04, 38/10, 38/16, 39/395, 45/00; C07K 14/46, 14/435, 16/30; C12N 5/20; G01N 33/53, 33/536

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

424/ 277.1; 435/7.1, 7.23, 240.27; 530/324, 327, 350, 387.7, 387.9, 388.8, 389.7, 391.1, 391.3

Form PCT/ISA/210 (extra sheet)(July 1992)*